

HERMES MEASUREMENTS OF THE GENERALIZED GDH-INTEGRAL AND OF QUARK-HADRON DUALITY

WOLF-DIETER NOWAK
DESY ZEUTHEN

ON BEHALF OF THE HERMES COLLABORATION

SPIN2002

BROOKHAVEN/USA, SEPT. 9, 2002

(1) GENERALIZED GERASIMOV-DRELL-HEARN INTEGRAL

- * MOTIVATION
- * ANALYSIS OF HERMES DATA
- * Q^2 -DEPENDENCE FOR DEUTERON, PROTON AND NEUTRON
 - NUCLEON RESONANCE REGION
 - DEEP INELASTIC SCATTERING (DIS) REGION
- * PROTON-NEUTRON DIFFERENCE

(2) QUARK-HADRON DUALITY

• REAL-PHOTON ABSORPTION BY A NUCLEUS:

[S.B. Gerasimov, Sov. J. Nucl. Phys. 2 (1966) 430]

[S.D. Drell, A.C. Hearn, Phys. Rev. Lett. 16 (1966) 908]

[H. Ahrenhoevel, nucl-th/0006083]

$$I_{GDH} = \int_{\nu_0}^{\infty} [\sigma^{\overleftarrow{}}(\nu) - \sigma^{\overrightarrow{}}(\nu)] \frac{d\nu}{\nu} = -\frac{4\pi^2 I \alpha}{M_t^2} \kappa^2$$

CONNECTS HELICITY-DEPENDENT DYNAMICS AND
STATIC GROUND-STATE PROPERTY OF NUCLEUS(ON)

$\sigma^{\overleftarrow{}}$, $\sigma^{\overrightarrow{}}$: photoabsorption cross sections for anti-parallel, parallel spin orientation

I, M_t, α : nucleus spin, mass, electromagnetic fine-structure constant

κ : anomalous contribution to magnetic moment of nucleus(on)

ν : photon energy in the target rest frame, ν_0 : photoabsorption threshold

target	κ	$I_{GDH}^{\gamma p}$ in μb
proton	1.79	-204
neutron	-1.91	-233
deuteron	-0.143	-0.65

• POLARIZED DIS ON NUCLEUS(ON)

⇒ GENERALIZED GERASIMOV-DRELL-HEARN INTEGRAL:

[R. Pantförder, hep-ph/9805434, and references therein]

$$I_{GDH}(Q^2) = \int_{\nu_0}^{\infty} [\sigma^{\overleftarrow{}}(\nu, Q^2) - \sigma^{\overrightarrow{}}(\nu, Q^2)] \frac{d\nu}{\nu}$$

NICE TOOL TO STUDY TRANSITION OF HELICITY-
DEPENDENT DYNAMICS FROM (PERTURBATIVE) DIS
REGION TO (NON-PERTURB.) PHOTOABSORPTION

$\sigma^{\overleftarrow{}}$, $\sigma^{\overrightarrow{}}$: virtual-photon absorption cross sections for anti-parallel, parallel spin orientation

Q^2 : photon virtuality

• SPIN $\frac{1}{2}$ TARGET:

$$I_{GDH}(Q^2) = \frac{8\pi^2\alpha}{M} \int_0^{x_0} \frac{g_1(x, Q^2) - \gamma^2 g_2(x, Q^2)}{K} \frac{dx}{x}$$

g_1, g_2 : polarised structure functions of the nucleon

$$\gamma^2 = Q^2/\nu^2, \quad x = Q^2/2M\nu, \quad x_0 = Q^2/2M\nu_0$$

K : virtual-photon flux factor

ν : photon energy in the target rest frame, ν_0 : photoabsorption threshold

FOR LARGER VIRTUALITY ($Q^2 \geq 1 \text{ GeV}^2$):

$$\int_0^1 g_2(x, Q^2) dx = 0 \quad (\text{BURKHARDT-COTTINGHAM SUM RULE})$$

$$I_{GDH}(Q^2) = \frac{16\pi^2\alpha}{Q^2} \Gamma_1(Q^2) \quad \text{with} \quad \Gamma_1(Q^2) = \int_0^1 g_1(x, Q^2) dx$$

GENERALIZED GDH INTEGRAL GIVEN BY FIRST
MOMENT OF SPIN STRUCTURE FUNCTION $g_1(x, Q^2)$

\implies DOES THE 'HARD $1/Q^2$ -REGIME'
STAY INTACT
DOWN TO LOWEST MEASURED Q^2 ?

- GDH p–n DIFFERENCE \longrightarrow BJORKEN SUM RULE:

$$I_{GDH}^p(Q^2) - I_{GDH}^n(Q^2) \xrightarrow{Q^2 \text{ large}} \frac{16\pi^2\alpha}{Q^2} [\Gamma_1^p(Q^2) - \Gamma_1^n(Q^2)]$$


FUNDAMENTAL BJORKEN SR DERIVED FROM CURRENT ALGEBRA AND ISOSPIN SYMMETRY:

$$\Gamma_1^p(Q^2) - \Gamma_1^n(Q^2) = \frac{1}{6} \cdot g_a \cdot C_{ns}(\alpha_s(Q^2))$$

$g_a = |g_A/g_V| = 1.2670 \pm 0.0035$: NEUTRON BETA-DECAY COUPLING CONSTANT

Q^2 -DEPENDENCE: GIVEN BY DGLAP-EVOLUTION

\Rightarrow GDH DATA FROM HERMES KINEMATICS
 ESSENTIAL TO CONNECT
 TWO IMPORTANT PROPERTIES OF THE NUCLEON:
 κ AND g_A



- Tracking: forward dipole magnet spectrometer with 57 chamber planes ($40 \text{ mrad} < \theta < 220 \text{ mrad}$); resolution: $\delta\theta < 0.6 \text{ mrad}$, $\delta p/p < 1.5\%$
- Particle ID: threshold/Ring-imaging(>1998) Cerenkov, TRD, preshower (H2), lead-glass calorimeter: e/h misidentification $< 0.4\%$
- Fast Trigger: scintillator hodoscopes H0, H1, H2; calorimeter: energy threshold 1.4 GeV

- REACTION:

DOUBLY POLARIZED INCLUSIVE e^+ OR e^- SCATT. ON ^2H (^1H)

- LONGITUDINALLY POLARIZED BEAM:

$E_0=27.6$ GeV, $\langle P_B \rangle = 0.55 \pm 0.01_{\text{syst.}}$ ($0.55 \pm 0.02_{\text{syst.}}$)

- LONGITUDINALLY POLARIZED INTERNAL GAS TARGET:

$\langle P_T \rangle = 0.85 \pm 0.03_{\text{syst.}}$ ($0.86 \pm 0.04_{\text{syst.}}$)

- INTEGRATED LUMINOSITY / NUCLEON: 111 (70) pb^{-1}

- KINEMATICS: $1.2 < Q^2 < 12$ GeV^2 in 6 bins

$$1.0 < W^2 < 45 \text{ GeV}^2$$

W : γ^* -nucleon invariant mass

\implies SUFFICIENT KINEMATIC COVERAGE AND STATISTICS TO STUDY FOR THE 1ST TIME THE GENERALIZED GDH INTEGRAL ON DEUTERON, PROTON AND NEUTRON SEPARATELY IN DIS REGION AND NUCLEON-RESONANCE REGION:

Region	W^2 -range	Statistics ^2H (^1H)
nucleon-resonance	$1.0 < W^2 < 4.2 \text{ GeV}^2$	0.13 M (0.55 M)
DIS	$4.2 < W^2 < 45 \text{ GeV}^2$	1.4 M (7.7 M)
unmeasured	$W^2 > 45 \text{ GeV}^2$	extrapolation [*]

* N. Bianchi, E. Thomas, Phys. Lett. B450 (1999), 439

Photon flux factor: $K = \nu \sqrt{1 + \gamma^2}$ (Gilman notation):

$$I_{GDH}(Q^2) = \frac{8\pi^2 \alpha}{Q^2} \int_0^{x_0} \frac{1}{\sqrt{1 + \gamma^2}} \tilde{A}_1 F_1 dx$$

$F_1 = F_2(1 + \gamma^2)/(2x(1 + R))$ and F_2 : unpolarized structure functions

F_2 by fit of A. Bodek, Phys. Rev. D8 (1973) 2331

$R = \sigma_L/\sigma_T$: ratio of absorption cross sections for long. and transverse virtual photons

(Resonance region: $R = 0.18$, DIS region: fit L. W. Whitlow et al., Phys. Lett. B 250 (1990) 193)

MEASURED LONGITUDINAL
CROSS-SECTION ASYMMETRY

PHOTON-HELICITY
ASYMMETRY

$$A_{\parallel} = \frac{1}{P_B P_T} \frac{N^{\leftarrow} L^{\rightarrow} - N^{\rightarrow} L^{\leftarrow}}{N^{\leftarrow} L^{\rightarrow} + N^{\rightarrow} L^{\leftarrow}} \quad \tilde{A}_1 = \frac{A_{\parallel}}{D} - \eta \tilde{A}_2$$

$N^{\leftarrow}, N^{\rightarrow}$: no. of scattered e^+ (e^-) for anti-parallel, parallel spin orientation

$L^{\leftarrow}, L^{\rightarrow}$: deadtime-corrected luminosity for anti-parallel, parallel spin orientation
 $y = \nu/E_0$: inelasticity of the reaction; $\eta(y, \gamma^2)$: kinematic factor

$D(y, Q^2, \gamma^2, R)$: virtual-photon (de)polarization factor

\tilde{A}_2^d, A_2^p : Resonance region: 0.06 (0.); DIS region: $c M x/\sqrt{Q^2}$ with $c=0.5$ (0.05)


FOR SPIN $\frac{1}{2}$ TARGET: $\tilde{A}_1 \equiv A_1$: VIRTUAL-PHOTON ASYMMETRY

• GENERALIZED GDH INTEGRAL FOR THE NEUTRON:

$$I_{GDH}^n = \frac{I_{GDH}^d}{1 - 1.5\omega_d} - I_{GDH}^p$$

I_{GDH}^n : C. Ciofi degli Atti et al., Phys. Lett. B 376 (1996) 309

$\omega_d = 0.058 \pm 0.010$: probability of the deuteron to be in a D-state
(M. Lacombe et al., Phys. Lett. B 101 (1981) 139)




RESONANCE CONTRIBUTION:
SMALL FOR $Q^2 > 3 \text{ GeV}^2$

Model using Q^2 -evolution of nucleon resonance amplitudes:

[I.G.Aznauryan, Phys. of At. Nuc., 58(1995)1014]

n-data: well described

p- and d-data: described within total exp. errors



DIS CONTRIBUTION: SIZEABLE DOWN TO LOWEST MEASURED Q^2 .


Mult.-Reggeon-exchange param. for $\sigma^{\leftarrow} - \sigma^{\rightarrow}$ at high energy

(\rightarrow extrap. into unmeas. region)

[N. Bianchi, E. Thomas, Phys. Lett. B450(1999), 439]

p-data: well described

d- and n-data: $Q^2 > 3 \text{ GeV}^2$: ok



DIS & RESONANCE REGION:

Model: leading-twist Q^2 -evol. of 1st moments of g_1, g_2


(w/o any resonance contribution)

[J.Soffer, O.V.Teryaev, Phys. Rev. D56(1997)7458]

\rightarrow all data well described

\Rightarrow hint on quark-hadron duality


• Q^2 -DEPENDENCE $\sim 1/Q^2$?



Dividing out $\sim 1/Q^2$ dependence (leading-twist) leads to flat Q^2 -behaviour for both p and n

\implies DOWN TO $Q^2 \simeq 1.5 \text{ GeV}^2$
 NO INDICATIONS FOR SIZEABLE CONTRIBUTIONS FROM HIGHER-TWISTS OR NUCLEON-RESONANCE FORM FACTORS

• PROTON-NEUTRON DIFFERENCE



• AT $Q^2 = 5 \text{ GeV}^2$: AGREEMENT WITH BJORKEN SUM RULE PREDICTION ($16.33 \pm 0.45 \mu\text{b}$) AND SLAC E-143/155 DATA)

• NOTE: Real-photon point $I_{GDH}^p|_{Q^2=0} - I_{GDH}^n|_{Q^2=0} = 29 \mu\text{b} \implies$ turn-over point not seen in HERMES kinematics

• Q^2 -dependence of p-n difference also follows $1/Q^2$ behaviour down to lowest measured values

• DEVELOPMENT OF DUALITY CONCEPT OVER HISTORY

* Origin from hadron-hadron scattering:

relationship between s - and t - channel behaviours of scatt. ampl.

[R. Dolen, D. Horn, C. Schmid, Phys. Rev. Lett. 19 (1967) 402; Phys. Rev. 166 (1968) 1768]

[G. Veneziano, Nuovo Cimento 57A (1968) 1395]

* Generalization of s - and t - channel duality:

including both resonant and non-resonant contrib.'s to cross section

[H. Harari, Phys. Rev. Lett. 20 (1968) 1395]

* In Inclusive electron scattering:

duality between resonances and valence quarks


(resp. background dual to sea quarks)

[W. Melnitchouk, hep-ph/0106262]

• BLOOM-GILMAN DUALITY: DIS \iff RESONANCE REGION

[E.D. Bloom, F.J. Gilman, Phys. Rev. Lett. 25 (1970) 1140; Phys. Rev. D4 (1971) 2901]

A STRUCTURE
 FUNCTION (OR A
 RATIO) MEASURED
 IN THE DIS REGION
 REPRESENTS AN
 AVERAGE OF THE
 SAME FUNCTION
 (OR RATIO)
 MEASURED IN THE
 RESONANCE REGION



• PQCD RE-FORMULATION OF BLOOM-GILMAN DUALITY:

- * Leading terms (non-interacting quarks) \rightarrow scaling
- * Non-leading terms ($q - g$ interactions) $\sim 1/Q^2$

[A. De Rujula, H. Georgi, H.D. Politzer, Phys. Lett. B64 (1976) 428]

DUALITY CONCEPT SUCCESSFUL IN THOSE RARE CASE(S) WHEN A PQCD CALCULATION DESCRIBES THE AVERAGE OF A HADRONIC OBSERVABLE (LIKE RES. REGION \leftrightarrow DIS)


but: pQCD picture must fail for low virtuality

\Rightarrow breakdown of Duality unavoidable at low Q^2

• POLARIZED ELECTRON-NUCLEON SCATTERING:

DUALITY IN *polarized* STRUCTURE FUNCTION (RATIO) IS *non-trivial* BECAUSE OF DIFFERENT HELICITY STRUCTURE OF POLARIZED DIS AND NUCLEON RESONANCE(S) [Δ : OPPOSITE HELICITY]

\Rightarrow INVESTIGATE Q^2 -DEPENDENCE OF $\Gamma_1^{res} / \Gamma_1^{DIS}$



1st TESTS IN POLARIZED CASE:

- **SLAC E143:** K. Abe et al., Phys. Rev. D 58, 112001 (1998)

- **HERMES KINEMATICS:**
DUALITY HOLDS OVER FULL MEASURED REGION
 \rightarrow same behaviour as known from unpolarized case

□ GENERALIZED GDH INTEGRALS FOR DEUTERON, PROTON AND NEUTRON MEASURED **FOR THE 1ST TIME MEASURED SIMULTANEOUSLY** IN NUCLEON-RESONANCE AND DIS REGION

▷ **RESONANCE CONTRIBUTION:** SMALL FOR $Q^2 > 3 \text{ GeV}^2$

▷ **DIS CONTRIBUTION:** SIZEABLE OVER FULL MEASURED Q^2

▷ **COMBINED DATA:** NO EFFECT FROM HIGHER-TWISTS OR RESONANCE FORM FACTORS DOWN TO $Q^2 \simeq 1.5 \text{ GeV}^2$

□ GDH PROTON-NEUTRON DIFFERENCE:

▷ **AGREES WELL WITH BJORKEN SUM RULE PREDICTION AND SLAC DATA AT $Q^2 = 5 \text{ GeV}^2$**

▷ **LEADING-TWIST $1/Q^2$ -BEHAVIOUR INTACT DOWN TO $Q^2 \simeq 1.5 \text{ GeV}^2$**

□ TURN-OVER OF PROTON AND DEUTERON GDH DATA (TOWARDS REAL-PHOTON POINT) NOT OBSERVED

□ **QUARK-HADRON DUALITY IN POLARIZED eN SCATTERING:**

⇒ **'HARD REGIME' (AGAIN) INTACT DOWN TO $Q^2 \simeq 1.5 \text{ GeV}^2$**